

Tryptic Soy Broth Irradiated

Art. No. 02-575

Also known as

Irradiated TSB

Specification

General liquid medium, sterilized by gamma-radiation to suit "media fill" tests in the pharmaceutical industry.

Formula* in g/L

Casein peptone.....	17,00
Soya peptone.....	3,00
Sodium chloride.....	5,00
Dipotassium phosphate.....	2,50
Dextrose.....	2,50
Final pH 7,3 ± 0,2 at 25°C	

* Adjusted and /or supplemented as required to meet performance criteria

Directions

Aseptically dissolve 30 g in 1 L of sterile distilled water and use as per standardized media fill run.

- For liquid filling tests:** Suspend 30 g in 1 L of sterile distilled water and mix well to dissolve. Use for liquid fill validation procedures.
- For solid filling tests:** Use powder for dry-fill validation procedures. Ensure that the final concentration of the medium is 30 g of Irradiated TSB suspended in 1 L of sterile distilled water.

Description

The Irradiated Tryptone Soy Broth (Irradiated TSB) is the classical TSB culture medium sterilized by a gamma-irradiation process making it appropriate for validation of aseptic filling processes in the pharmaceutical industry (Media-Fill Test).

The sterility of the dehydrated culture medium is verified according to the methodology as described in the pharmacopoeia. It is subjected to exactly the same conditions as the pharmaceutical product, including filling and closing to ensure that there is no microbial contamination occurring during the process.

The gamma-irradiation treatment is carried out according to the Annex B of the ISO 11137 Standard, and warrants a 25 kGy absorption by the medium, which is enough to suppress any vegetative and/or spore from of microorganisms present in the powder without modification of the medium's performance. All batches of Irradiated TSB are checked for sterility and performance.

Technique

All the conditions and data for the validation of aseptic filling process can be consulted in the ISO Standard 13408 -1:1998 in the chapters devoted to the methods of preparation of sterile products in several Pharmacopoeias.

References

- EUROPEAN PHARMACOPOEIA 7.0 (2011) 7th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- ISO Standard 11137:1995. Sterilization of health care products- Requirements for validation and routine control - Radiation Sterilization. (Annex B).
- ISO/TS 11133-1: 2009 Microbiology of food and animal feeding stuffs.- Guidelines on preparation and production of culture media. Part 1: General guidelines on quality assurance for the preparation of culture media in the laboratory.
- ISO/TS 11133-2: 2003 Corr. 2004 Microbiology of food and animal feeding stuffs.- Guidelines on preparation and production of culture media. Part 2: Practical guidelines on performance testing of culture media.
- ISO Standard 13408-1:1998. Aseptic processing of health care products-Part 1: General requirements.
- US PHARMACOPOEIA 28 / NATIONAL FORMULARY 23 (2005) General Chapters § <71> Sterility Tests y § <1208> Sterilization and Sterility Assurance of Compendial Articles.

Storage

For laboratory use only. Keep tightly closed, away from bright light, in a cool dry place (+4°C to 30°C and <60% RH).

(continues on the next page)

Tryptic Soy Broth Irradiated

Art. No. 02-575

T

Quality control

Incubation temperature: 35°C ± 2.0

Incubation time: 24 h-5 days

Inoculum: 10-100 CFU (according to standard ISO/TR 11133-1:2000 and ISO/TS 11133-2:2003)

Microorganism	Growth	Remarks
<i>Bacillus subtilis</i> ATCC 6633	Good	-
<i>Staphylococcus aureus</i> ATCC 6538	Good	-
<i>Clostridium sporogenes</i> ATCC 19404	Good	In anaerobic conditions
<i>Escherichia coli</i> ATCC 8739	Good	-
<i>Salmonella abony</i> NCTC 6017	Good	-
<i>Candida albicans</i> ATCC 10231	Good	-
<i>Aspergillus brasiliensis</i> ATCC 16404	Good	Black sporulation at 5 days



Left: *Escherichia coli* ATCC 8739
Centre: *Salmonella* sp. NCTC 6017
Right: Uninoculated tube (Control)



Left: Uninoculated Tube (Control)
Right: *Clostridium sporogenes* ATCC 19404